



CONSTRUCTING KNOWLEDGE



The König Group is composed by companies who have been in business within the veterinary and environmental health markets for more than 35 years.

Their business insight, expansive regional policy, sound commercial practices, quality of their products, constant innovations in new developments and, above all, their human resources have enabled the **König Group** to warrant growth during all these years.



_MISSION

To meet and fulfill the needs and expectations of our Customers, by producing high quality products, using technology and processes that protect the environment and respect ethical and moral values in our industrial and commercial practices.

_VISSION

To consolidate its position at an international level and as a laboratory manufacturing high-quality product.

_OUR MAIN CORPORATE VALUES ARE SUMMARIZED BELOW:

Our people and teamwork.

Continuous innovation in processes and products.


Quality.

Flexibility in adaptation within unknown scenarios.

Solidarity and environmental care.



KONIG GROUP



PLANTS AND WAREHOUSES

Sao Paulo Plant: 2700 M2

Production of vitamin cores, antibiotics, coccidicides, nutritional improvement products and mineral concentrates.

Address: Rodovia Castelo Branco Km 68.5, Mairinque, Sao Paulo, Brazil

Avellaneda Logistics Center: 3300 m2

Stock of finished products and raw materials.

Address: Belisario Hueyo 210, Avellaneda, Buenos Aires, Argentina

Konig Plant: 2000 m2

Injectable products, tablets, hormonal products, antiparasitics, anesthetics, antibiotics and mineral supplements

Address: Av. Rivadavia 719, Avellaneda, Buenos Aires, Argentina.

Chivilcoy Plant, Buenos Aires Province, Argentina: 1850 m2

Production of Sprays

Address: Route 5, Km 160, Industrial Park, Chivilcoy, Bs. As., Argentina

Binka Plant: 1800 m2

Production of pesticides and healing substances.

Address: Méjico 1060, Pineyro, Bs. As., Argentina

Uruguayan Project

An investment of U\$S 4.600.000 has been planned for the year 2014 in order to build a 3000 m2 plant, to be located on a 9-hectares plot near the Carrasco Airport, at Montevideo, Uruguay.

The plant will be destined to the production of large volumes of injectable and oral antiparasitic products for cattle, sheep and pigs.

PLANTS AND WAREHOUSES

SAO PAULO PLANT : 2700 M2

Production of vitamin cores, antibiotics, coccidicides, nutritional improvement products and mineral concentrates.

Address: Rodovia Castelo Branco Km 68.5, Mairinque, Sao Paulo, Brazil



PLANTS AND WAREHOUSES

Chivilcoy Plant, Buenos Aires Province, Argentina: 1850 m2

Production of Sprays

Address: Route 5, Km 160, Industrial Park, Chivilcoy, Bs. As., Argentina



PLANTS AND WAREHOUSES

Uruguayan Project

An investment of U\$S 4.600.000 has been planned for the year 2014 in order to build a 3000 m² plant, to be located on a 9-hectares plot near the Carrasco Airport, at Montevideo, Uruguay.

The plant will be destined to the production of large volumes of injectable and oral antiparasitic products for cattle, sheep and pigs.





König Argentina

König Brasil

König Uruguay

Bolivia

Ecuador

El Salvador

Guatemala

Honduras

Méjico

Panamá

Paraguay

Perú

República Dominicana

Venezuela



PRODUCTS

LARGE ANIMALS



—
WWW.KONIGLAB.COM

ANTIINFLAMMATORY

ANALGESICS

WWW.KONIGLAB.COM

APLONAL 5%

Maximum potency non-steroid antiinflammatory, analgesic and antipyretic agent.



FOR USE IN



DISTRIBUTION COUNTRIES



Injectable sedative, myorelaxant, analgesic and anesthetic agent for small and large animals.

FORMULA

Each ml contains: xylazine hydrochloride 20 mg;
excipients and vehicle q.s.

INDICATIONS

- **Horses:** Colic and inflammatory injuries of the locomotive system.
 - **Cattle:** Supportive treatment in acute infectious diseases, calf diarrhea and fallen cow syndrome.
 - **Pigs:** Treatment of the metritis-mastitis-agalactia syndrome.
- Post-surgical and post-traumatic relief in all species

ADMINISTRATION AND DOSAGE

- **Horses:** Apply 1 ml/45 kg weight (1.1 mg flunixin/kg) by IM or IV route, every 24 hours up to a maximum of 5 consecutive days. Use the IV route for colics; the treatment can be repeated up to 3 times at 30 minute intervals.
- **Cattle:** Apply 2 ml/45 kg weight (2.2 mg flunixin/kg) up to a maximum of 5 days. For neonate diarrhea, 2 to 3 ml/calf, 1 to 2 applications.
- **Pigs:** Apply 2 ml/45 kg weight (2.2 mg flunixin/kg), every 24 hours during 3 days.

ANESTHESIC

MYORELAXANTS

WWW.KONIGLAB.COM

KENSOL

Myorelaxant and anesthetic sedative.



FORMULA

Each ml contains: xylazine hydrochloride 20 mg;
excipients and vehicle q.s.

INDICATIONS

Sedative, myorelaxant, analgesic and anesthetic
for use alone or in association with preanesthetics
or anesthetics for contention maneuvers or major
or minor surgery in all domestic animals.



FOR USE IN



DISTRIBUTION COUNTRIES



SEDOMIN

Sedative, myorelaxant, analgesic and anesthetic agent for horses and cattle. Concentrated dose.



Injectable sedative, myorelaxant, analgesic and anesthetic agent for horses and cattle.

FORMULA

Each ml contains: xylazine (as hydrochloride) 100 mg, excipients and vehicle q.s.

INDICATIONS

Diagnostic procedures, presurgical sedation, orthopedic and odontological procedures and minor surgery in horses. Transportation, minor and major surgery in cattle.

ADMINISTRATION AND DOSAGE

■ **Horses:** 1 ml/100 kg weight, via IV or 2 ml /100 kg weight, via IM. Administration of the indicated dose causes sedation for a period of 1 to 2 hours, with analgesia during the first 15 to 30 minutes.

■ **Cattle:** 0.05 - 0.1 ml/100 kg weight, via IM or IV for minor surgery, sedation or transportation. Onset of action at 30 seconds and lasts, approximately, 2 hours. For major surgery, 0.2 ml/100 kg, via IM or IV; duration of effect 2 to 4 hours.



FOR USE IN



DISTRIBUTION COUNTRIES



Sedomin (0.1 ml/100 kg) can also be combined with Vetanarcol (10 ml/100 kg). As complement of local anesthesia, **Sedomin** is used at the same dose indicated for sedation, concurrent with locally applied procaine or lidocaine. As preanesthetic for general anesthesia, **Sedomin** is used at the same dose indicated for sedation. Due to the depression additive effect on the central nervous system, only 1/2 to 1/3 of the barbituric dose must be administered to achieve an adequate anesthesia.

VETANARCOL

Safe anesthetic. Cardiovascular stimulant. No effects on respiratory system.



Dissociation acting general anesthetic for domestic and wild animals. Ideal for emergencies and traumatisms.

FORMULA

Each ml contains: ketamine (as hydrochloride) 50 mg,
excipients and vehicle q.s.

INDICATIONS

Vetanarcol is an anesthetic indicated for premedication, induction or maintenance use, alone or associated with major and minor tranquilizers, sedatives, narcotics, analgesics and other anesthetics for all kinds of domestic and wild animals.



FOR USE IN



DISTRIBUTION COUNTRIES



DERMATOLOGIC ANTIBIOTICS

& ANTIINFLAMMATORY PRODUCTS

WWW.KONIGLAB.COM

BTK 35

Cure and prevention of keratoconjunctivitis. Total acting bactericide, fungicide and viricide. Antiinflammatory.



Topical administration antibacterial and antiinflammatory agent for small and large animals.

FORMULA

Each 100 g of concentrate: argentic sulfadiazine 0,85 g%;
Oxytetracycline hydrochloride 5,00 g%; hydrocortisone
acetate 0.29 g%; excipients q.s.f.

INDICATIONS

For the prevention and treatment of infectious
keratoconjunctivitis in cattle, as well as for the prevention
and treatment of bacterial inflammation and infection
of traumatic and surgical wounds in horses.



FOR USE IN



DISTRIBUTION COUNTRIES



KURAMICINA L.A

Sustained action. Five-day bioavailability. Just one application is enough.



FOR USE IN



DISTRIBUTION COUNTRIES



Immediate and long acting injectable antibiotic.

FORMULA

Each 100 ml contains: amphoteric base oxytetracycline 20 g, excipients and vehicle q.s

INDICATIONS

For the prevention and cure of general infectious diseases, caused by Gram positive and Gram negative bacteria, micoplasmas, rickettsias and clamydias in cattle, sheep and pigs. Very effective in keratoconjunctivitis (intrapalpebral) and for the treatment of campylobacteriosis in bulls.

ADMINISTRATION AND DOSAGE

Apply 1 ml per each 10 kg weight, exclusively by IM. In less than 10 kg suckling pigs, apply a total of 1 ml, via IM or SC. Both for preventive and curing treatments, the frequency is just one application except if the disease is sufficiently acute to justify a second application within 3 to 5 days.

■ **Precautions:** According to the volume to be administered, the application must be performed in two or more injection sites, never exceeding the following volumes in each site: 20 ml in cattle, 5 ml in sheep and 10 ml in pigs.

ANTIDIARRHEIC PRODUCTS

WWW.KONIGLAB.COM

DIAFIN N

A single application stops diarrhea. Eliminates pathogen enterobacteria.



FOR USE IN



+

DISTRIBUTION COUNTRIES



Injectable intestinal motility modulator antibiotic and antidiarrheal for small animals.

FORMULA

Each ml contains benzetimide hydrochloride 0.165 mg;
enrofloxacin 25 mg; excipients and vehicle q.s.

INDICATIONS

Treatment of complicated bacterial or viral diarrheas
in dogs and cats.

ADMINISTRATION AND DOSAGE

Apply 1 ml per each 10 kg weight, via SC or IM to both species. In most cases, a single application is sufficient to cure diarrhea. Treatment can be repeated at 24 or 48 hours.

EXTERNAL

ANTIPARASITICS

WWW.KONIGLAB.COM

BACTROFLAY POUR ON

External Antiparasitic. Excellent Residual Power



FOR USE IN



DISTRIBUTION COUNTRIES



External pour-on antiparasitic without restriction as regards milk and minimum restriction as regards beef. Excellent residual power.

FORMULA

Contents per 100 ml. Ethion 15 grams;
Cypermethrin 5 grams; Excipient and Vehicle q.s.

INDICATIONS

Control and treatment of infestations by Haematobians
irritans (horn fly) in bovines.

ADMINISTRATION AND DOSAGE

Apply along the dorsal midline from withers to rump. The interval between applications should be 50 days.

■ **Calves weighing:** 100 to 200 K.: 5 mL. // 200 to 400 K.: 10 mL. // over 400 K.: 20 mL.

WAITING TIME:

■ **Milk Bovines:** 6 hours

■ **Beef Bovines:** 6 days

The restriction applies because of Cypermethrin, in fulfillment of Senasa Resolution N° 1343/93

BACTROFLY TAGS

External Antiparasitic. Excellent Residual Power



FOR USE IN



DISTRIBUTION COUNTRIES



External antiparasitic without restriction as regards beef and 48 hours as regards milking period.

FORMULA

Ethion 40g %, Excipients q.s. 100 g.

INDICATIONS

Control and treatment of infestation due to horn fly
in beef and milk bovines.

ADMINISTRATION AND DOSAGE

Place one tag (caravana) on each animal. Replace the tags every 4 months, once this period has been completed, removal is recommended in order to avoid the possible appearance of resistant populations because the flies are being exposed to sub-lethal doses. Place the tags using an applicator pincers that should be disinfected with an antiseptic solution before being used.

WAITING TIME:

No restriction in beef cattle.

Milking waiting period in Argentina: 48 hours.

INTERNAL

ANTIPARASITICS

WWW.KONIGLAB.COM

RICOVERN

Wide spectrum antiparasitic: adulticide, larvicidal and ovicidal. No irritating effect. Maximum bioavailability.



Internal, injectable, wide spectrum antiparasitic for bovines
(Wide spectrum gastrointestinal and lung antiparasitic).

FORMULA

Contents per each 100 mL: Ricobendazol 15 grams;
excipient and vehicle q.s.

INDICATIONS

Indicated for the treatment of infestations by nematodes:
Haemonchus spp., Trichostrongylus spp., Strongylus spp.,
Ostertagia spp., Nematodirus spp., Cooperia spp.,
Bunostomum spp. and Dictyocaulus spp. in cattle.

ADMINISTRATION AND DOSE

Apply subcutaneously in front of, or behind, the shoulder. The recommended dose is 1 mL for each 40 K. of weight.
Do not inject more than 10 mL per point of application.

RESTRICTION PERIOD

- **Milk cattle:** 3 days
- **Beef producing cattle:** 14 days.



FOR USE IN



DISTRIBUTION COUNTRIES



VERMIPLEX PLUS 500

Broad spectrum antiparasitic for horses, pregnant mares, Racing purebred horses in training.



FOR USE IN



DISTRIBUTION COUNTRIES



Total spectrum paste anthelmintic for horses with ovicide, larvicide and adulticide action.

FORMULA

Each 100 g contains: mebendazole 12.5 g, closantel 11.8 g
and excipients, q.s.

INDICATIONS

Treatment of Parasitosis due to *Gasterophilus* spp., *Parascaris equorum*, *Oxyuris equi*, *Habronema muscae*, *Trichostrongylus axei*, *Trichonema* spp., *Tridontophorus* spp., *Strongylus* spp., *Strongyloides westeri* and *Fasciola hepatica* in horses.

ADMINISTRATION AND DOSAGE

Oral administration of 26 g (one syringe)/500 kg weight, every three months. This drug can be administered to pregnant mares and foals.

PHYSIOLOGICAL CICATRIZATION PRODUCTS

WWW.KONIGLAB.COM

KURADERM PLATA

Physiological Healing Agent. Keratoplastic. No formation of crust. Accidental or surgical wounds.



Antimicrobial; healing; astringent and high adherence hemostatic agent, for topical use in small and large animals.

FORMULA

100 g of concentrate contains: Argentic sulfadiazine 1.25 g;
neomycin base (as sulfate) 5.4 g; aluminum 8.33 g
and excipients q.s. Propellant: Butane-propane.

INDICATIONS

Prevention and treatment of infectious processes in wounds.
Hemostatic effect. Healing of wounds, eschars,
burns and ulcers.

ADMINISTRATION AND DOSAGE

Shake before applying. After cleansing the area to be treated, spray a thin layer on same from a distance of approximately 10 cm ensuring penetration inside the wound. Repeat once a day until the wound heals.



FOR USE IN



DISTRIBUTION COUNTRIES



BACTROVET PLATA AM

High adherence hemostatic, healing and antimyasic agent.



Topical use, high adherence, healing, antimyasic, antimicrobial and hemostatic agent.

FORMULA

Each 100 g of concentrate contains: Argentic sulfadiazine 0.1 g; aluminum 5 g; DDVP 1.6 g; cypermethrin 0.4 g and excipients q.s.

INDICATIONS

Prevention of bleeding, infection and myiasis in traumatic and surgical wounds. Curing treatment of ulcerous skin myiasis, with larvicide and healing stimulant effect.

For use in small and large animals.

ADMINISTRATION AND DOSAGE

Shake before using. Spray abundantly on the wound or myiasis at an approximate distance of 10 cm, ensuring penetration inside the wound. Apply once daily until healing.



FOR USE IN



DISTRIBUTION COUNTRIES



BACTROVET PLATA PASTA

Intensive high-efficacy de-wormer Paste



FOR USE IN



DISTRIBUTION COUNTRIES



Antimiasic, repellent, bactericide, antifungal, wound healing, epithelization and hemostatic agent.

FORMULA

Contents per each 100 grams: silver sulphadiazine. 0,1 grams;
Aluminium, 5,0 grams; Imidacloprid 0,7 grams; cypermethrine
3,0 grams; excipient q.s.

INDICATIONS

Treatment of any type of wound in any species. Treatment and prevention of worms (bicheras) in surgical interventions such as castration, dehorning, tail docking, navels of calves, cracks in hooves and wounds of any origin. Wounds in general.

ADMINISTRATION AND DOSAGE

Place the paste in layers on the area to be treated, ensuring penetration of the product into the wound. Also apply around the wound, forming a 5-cm. halo. The number of applications depends on the type of wound and environmental conditions. The silver color indicates that the animal has been treated, The number of applications, as preventive action, depends on the type of wound and environmental conditions, but in the case of navel worms (bichera) . a single application is recommended within 24 hours after birth. In the case of castration wounds: 2 separate applications with an interval of 72 hours are recommended.

KOBRA

High adherence, healing and antimyasic agent.



Topical use, high adherence, healing, antimyasic agent.

FORMULA

Each 100 g of concentrate contains: DDVP, 1,26 g;
Cypermethrin, 1,30 g and excipients q.s.

INDICATIONS

Prevention of bleeding, infection and myiasis in traumatic and surgical wounds. Curing treatment of ulcerous skin myiasis, with larvicide and healing stimulant effect.

For use in small and large animals.

ADMINISTRATION AND DOSAGE

Shake before using. Spray abundantly on the wound or myiasis at an approximate distance of 10-15cm, ensuring penetration inside the wound. Apply once daily until healing. Recommended in navel, an application within 24 hours of birth, in castration wound two separate applications for 72 hours.



FOR USE IN



DISTRIBUTION COUNTRIES



ENDOECTOPARASITICIDES

WWW.KONIGLAB.COM

AVOMETIN + ADE

Vitamin supplement for bovines.



FOR USE IN



DISTRIBUTION COUNTRIES



Internal parasiticide, injectable vitamin supplement for bovines.

FORMULA

Abamectin, 1.1 g; vitamin A (palmitate), 5,000,000 UI;
vitamin D3, 40,000 UI; vitamin E (acetate), 1,000 UI;
excipients q.s.f.; 100 mL.

INDICATIONS

External and Internal parasiticide. (*)

■ **INTERNAL PARASITES: Gastrointestinal parasites:** Ostertagia ostertagi (including inhibited stage), Haemonchus placei, Mecistocirrus digitatus, Trichostrongylus axei, Trichostrongylus colubriformis, Cooperia oncophora, Cooperia punctata, Cooperia pectinata, Cooperia spp., Oesophagostomum radiatum, Nematodirus helvetianus (adults), Nematodirus spathiger, Strongiloides papillosus, Bunostomum phlebotomum; Toxocara vitolorum. **Pulmonary parasites:** Dictyocaulus viviparus (including inhibited stages).
■ **EXTERNAL PARASITES:** Dermatobia hominis; Hypoderma spp., Linognathus vituli; Haematophinus eurysternus; Solenopotes capillatus; Psoroptes bovis; Sarcoptes scabiei var. bovis; Damalinia bovis; Ornithodoros savignyi; Boophilus microplus (apply every 21 days), Boophilus decoloratus; Chorioptes bovis. To prevent and treat vitamin A, E and D3 deficiencies.

ADMINISTRATION AND DOSAGE

1 ml for every 50 kg of live weight, by subcutaneous route.

WITHHOLDING PERIOD

35 days. Do not administer to dairy cows in production.

RANDON

Parasticide external and internal.



Injectable endectocide for bovines, sheep and pigs.

FORMULA

Ivermectin, 1.00 g%; excipients q.s.f., 100 mL.

INDICATIONS

Internal and external parasiticide. (*)

(*) **Bovines, sheep and porcines:** gastrointestinal parasites (*Ostertagia ostertagi*, *Haemonchus* spp., *Trichostrongylus axei*, *Cooperia oncophora*, *Cooperia punctata*, *Cooperia pectinata*, *Oesophagostonun* spp.); pulmonary parasites (*Dictyocaulus viviparus* y *filaria*); mites (*Psoroptes bovis*); biting lice and sucking lice (*Linognathus vituli*; *Haematophinus eurysternus*; *Solenopotes capillatus*) and larvas causing myiasis (*Dermatobia hominis*).

ADMINISTRATION AND DOSAGE

Subcutaneous application.

■ **Bovines and sheep:** 1 mL every 50 kg of weight.

■ **Porcines:** 1 mL every 33 kg of weight.

WITHHOLDING PERIOD

Restriction period in animals destined for human consumption: **Bovines:** 35 days. **Sheep:** 21 days. **Porcines:** 18 days. In dairy cows do not apply within 28 days before parturition. Do not apply in bovines and sheep in production whose milk is destined for human consumption.



FOR USE IN



DISTRIBUTION COUNTRIES



RANDON + ADE

Internal parasiticide and injectable vitamin supplement.



FOR USE IN



DISTRIBUTION COUNTRIES



Internal parasiticide and injectable vitamin supplement for bovines, sheep, goats, horses and pigs.

FORMULA

Ivermectin, 1.1 g%; vitamin A (palmitato), 5,000,000 UI,
vitamin D3, 40,000 UI; vitamin E (acetate), 1,000 UI;
excipients q.s.f., 100 mL.

INDICATIONS

Internal and external parasiticide. (*)

(*) **Bovines, sheep and pigs:** gastrointestinal parasites (*Ostertagia ostertagi*, *Haemonchus* spp., *Trichostrongylus axei*, *Cooperia oncophora*, *Cooperia punctata*, *Cooperia pectinata*, *Oesophagostomum* spp.); pulmonary parasites (*Dictyocaulus viviparus* y *filaria*); mites (*Psoroptes bovis*); biting lice and sucking lice (*Linognathus vituli*; *Haematophinus eurysternus*; *Solenopotes capillatus*) and larvae causing myiasis (*Dermatobia hominis*). Multivitamin: to prevent and treat deficiencies in vitamins A, E and D3.

ADMINISTRATION AND DOSAGE (subcutaneous application)

■ **Bovines, sheep, goats and horses:** 1 mL every 50 kg of weight.

■ **Porcines:** 1 mL every 33 kg of weight.

WITHHOLDING PERIOD

Restriction period in animals destined for human consumption: 1 month. In dairy cows do not apply within 28 days before parturition. Do not apply in bovines and sheep in production whose milk is destined for human consumption.

RANDON TITANIUM

Internal and external parasiticide



FORMULA

Ivermectin 3,15 %; excipients q.s.f. 100 mL

INDICATIONS

Endoparasiticide: Gastrointestinal and pulmonary nematodes (adults and L4). Ectoparasiticide: Ticks *Bombus microplus*; Botfly *Dermatobia hominis*; *Cochliomya hominivora*; Lice *Haematopinus*, *Linognathus*, *Solenopotes*; Scabies *Psoroptes*, *Sarcoptes*.

ADMINISTRATION AND DOSAGE

Bovines: Subcutaneous injection 1 mL/50 kg bw.

Shake well for a minute before use, and keep shaking frequently during treatment

WITHHOLDING PERIOD

4 months



FOR USE IN



DISTRIBUTION COUNTRIES



HORMONAL PRODUCTS

WWW.KONIGLAB.COM

ECP ESTRADIOL

Minimum dose and long acting estrogen.



+

FOR USE IN



+

DISTRIBUTION COUNTRIES



Injectable hormonal agent for small and large animals.

FORMULA

Each 100 ml contains: estradiol cipionate 0.2 g;
excipients and vehicle q.s.

INDICATIONS

■ **Cows and heifers:** Anestrus without follicular cyst; retention of the corpus luteum in milk cows; expulsion of placental residues or mummified foetus. Mares, ewes and sows: Anestrus.

ADMINISTRATION AND DOSAGE (IM Injection only.)

■ **Cows:** anestrus: 1.5 to 2.5 ml total dose; corpus luteum persistence: 2.0 ml total dose; amnion hydrops: 3.0 to 4.0 ml total dose; pyometra, placenta retention and expulsion of mummified foetus: 5.0 ml total dose. Heifers: Anestrus: 1.5 ml total dose; Mares: anestrus: 2.5 to 5.0 ml total dose; Ewes and sows: Anestrus: 0.25 to 0.5 ml total dose.

MINERALIZING PRODUCTS

WWW.KONIGLAB.COM

ENERGO-MAG

Prevention and Control of hypomagnesemia.



Injectable mineral supplement for sheep and cattle.

FORMULA

Each 100 ml contains: magnesium

5-oxo-2-pyrrolidinecarboxylate 60 g; excipients and vehicle q.s.

INDICATIONS

Prevention agent in rodeos or flock of sheep and in animals with a history of hypoenergetic and hypomagnesemia syndrome. (*)

(*) Application recommended in strongly stressing conditions such as: transportation, weaning, storms, winter grazing, shearing of sheep, initiation of the last third of pregnancy in sheep. Elected therapeutic agent in the treatment of total metabolic dysfunction originated by a sudden alteration of energetic and/or magnesium requirements in ruminants: grass tetany; milk cow ketosis; fat cow syndrome; pregnancy toxemia in sheep; hypoenergetic syndrome in sheep.

ADMINISTRATION AND DOSAGE

Preventive dose: apply 1 to 1.5 ml/10 kg weight by SC route. Curing dose: 2.5 ml/10 kg weight, divided into equal parts, administering half of the dose by slow IV and the other half by SC.



FOR USE IN



DISTRIBUTION COUNTRIES



GLYPONDIN

Organic modifier; ideal for fattening and weaning. Immunostimulant. Fertility enhancement.



Sustained release mineral injectable supplement for cattle. 120 day duration.

FORMULA

Each 100 ml contains: copper as calcium cupric ethylenedinitrilotetraacetate 2.5 g, excipients and vehicle q.s. (25 mg active copper per ml).

INDICATIONS

Treatment of copper deficient conditions in cattle. Prevention and curing agent in hypocuprosis endemic zones.

ADMINISTRATION AND DOSAGE

SC exclusive administration of 1 ml/100 kg weight. Guidance: cows, bulls, young bulls and heifers, 4 ml; calves of over 200 kg weight: 3 ml. Calves of up to 200 kg weight: 2 ml. Repeat application every 3 to 4 months.



FOR USE IN



DISTRIBUTION COUNTRIES



GLYPONDÍN 4 MULTIMINERAL

Injectable Multiminerall Supplement.



FOR USE IN



DISTRIBUTION COUNTRIES



Multiminerall Supplement containing Zinc, Copper, Manganese and Selenium, for cattle.

FORMULA

Contents per each 100 mL: EDTA Disodium Manganese 7,10 grams; EDTA Disodium Zinc 24,45 grams; EDTA Disodium Copper 12,50 grams; Sodium Selenite, 1,10 grams; excipient q.s.

INDICATIONS

Preventive and healing in cases of zinc, copper, manganese and selenium deficiencies in bovines. Optimises production, reproductive function and immune system.

ADMINISTRATION AND DOSAGE

Administer exclusively via subcutaneous route.

Animals up to one year old: 1 mL per every 50 Kg. live weight (equivalent to zinc: 0.80 mg/Kg.; copper: 0.40 mg/Kg.; manganese: 0.20 mg/Kg.; selenium: 0.10 mg/Kg.)

Animals from one to two years old: 1 mL per every 75 Kg. live weight (equivalent to zinc: 0.53 mg/Kg.; copper: 0.265 mg/Kg.; manganese: 0.13 mg/Kg.; selenium: 0.06 mg/Kg.)

Animals over 2 years old: 1 mL per each 100 Kg. live weight (equivalent to zinc: 0.40 mg/Kg.; copper: 0.20 mg/Kg.; manganese: 0.10 mg/Kg.; selenium: 0.05 mg/Kg.)

REVIVON

Mineral supplement for bovines, horses, sheep and pigs.



FOR USE IN



DISTRIBUTION COUNTRIES



Glucose restorative and injectable mineral supplement for bovines, horses, sheep and pigs.

FORMULA

Every 100 mL contains: anhidre glucose, 10.000 g%;
vitamin B1, 0.100 g%; vitamin B2, 0.010 g%;
nicotinic acid, 0.100 g%; sodium chloride, 0.514 g%;
potassium chloride, 0.089 g%; calcium chloride, 0.028 g%;
magnesium chloride, 0.014 g%; caffein, 0.400 g%;
excipients q.s.f., 100 mL.

INDICATIONS

Treatment for acetonemia; downer cow syndrome, weakening,
base treatment for mineral shortages; rise in yield
of dairy cows, recovery due to energy loss in racing horses.

ADMINISTRATION AND DOSAGE (by subcutaneous injection, intramuscular or intravenous)

■ **Bovines and horses:** 1 mL per kg of live weight.

■ **Sheep and pigs:** 5 mL per kg of live weight.

VITA 3

Mineral and vitamin supplement for bovines.



Injectable mineral and vitamin supplement for bovines.

FORMULA

Every 100 mL contains: vitamin A (palmitate), 5,000,000 UI;
vitamin D3, 2,000,000 UI; vitamin E, 1,000 UI;
calcium borogluconate, 10,000 mg;
magnesium hypophosphite, 7,000 mg; sodium selenite,
150 mg; excipients s.q.f. 100 mL.

INDICATIONS

To improve mineral and vitamin deficiencies. General booster.

ADMINISTRATION AND DOSAGE (*subcutaneous application*)

5 to 10 mL per animal.



FOR USE IN



DISTRIBUTION COUNTRIES



GASOL EXTRA FORTE

Injectable intraruminal for bovines.



Anti bloat agent (Carminative) injectable intraruminal for bovines.

FORMULA

Poloxalene, 94 g%; excipients q.s.f., 100 g.

INDICATIONS

For the prevention and treatment of meteorosis in cattle, as a result of ingestion of legumes (alfalfa, clover, etc.).

ADMINISTRATION AND DOSAGE *(intraruminal application)*

- **Preventive treatment:** 10 mL for every 450 kg of live weight. Administer 24 hours before introducing animals to high-risk pastures; complete prevention with plenty of drinking water and fibrous foodstuff.
- **Curative treatment:** 25 mL every 450 kg of live weight.



FOR USE IN



DISTRIBUTION COUNTRIES



VITATONE

Mineral and vitamin supplement for bovines.



Injectable mineral and vitamin supplement for bovines.

FORMULA

Vitamin A (palmitate), 2,000,000 UI; vitamin D3, 2,000,000 UI;
vitamin E, 1,000 UI; copper sulphate, 1,000 mg; zinc sulphate,
80 mg; cobalt chloride, 60 mg; sodium selenite, 15 mg;
excipients s.q.f. 100 mL.

INDICATIONS

Vitamin and mineral booster for development in bovines
especially calves. Stimulant for organic defenses.

ADMINISTRATION AND DOSAGE *(subcutaneous application)*

- **Calves:** 10 mL per animal.
- **Adults:** 10 to 20 mL per animal



FOR USE IN



DISTRIBUTION COUNTRIES



PRODUCTS

SMALL ANIMALS



—

WWW.KONIGLAB.COM

SMALL ANIMALS

ANALGESICS

ANTIINFLAMMATORY PRODUCTS

WWW.KONIGLAB.COM

APLONAL 1%

Maximum potency non-steroid antiinflammatory, analgesic and antipyretic agent.



Injectable non-steroid antiinflammatory, analgesic and antipyretic agent for dogs,
with a morphine equivalent potency.

FORMULA

Each ml contains: base flunixin (as meglumine) 10 mg;
excipients and vehicle q.s.

INDICATIONS

Pain and inflammation relief in musculo-skeletal disorders;
endotoxic shock; post-traumatic and post-surgery pain.

■ **In ophthalmology:** surgery, uveitis and conjunctivitis.

Supportive treatment of parvovirus in dogs.

ADMINISTRATION AND DOSAGE (via IM or IV)

Apply 1 ml per 10 kg weight, via IM or IV, every 24 hours. Maximum 3 applications.



FOR USE IN



DISTRIBUTION COUNTRIES



APLONAL 20mg

The maximum potency, non-steroid anti-inflammatory, analgesic and antipyretic agent for ambulatory treatment.



FOR USE IN



DISTRIBUTION COUNTRIES



A non-steroid anti-inflammatory, analgesic and antipyretic agent for oral administration in dogs with a potency equivalent to morphine.

FORMULA

Each tablet contains: flunixin base (as meglumine) 20 mg
and excipients, q.s.p. 300 mg.

INDICATIONS

Pain and inflammation relief in dogs, irrespective of its origin,
especially musculo-skeletal.

ADMINISTRATION AND DOSAGE

1 tablet per 20 Kg weight every 24 hours, preferably with food, during a maximum of 3 days.

In chronic cases alternate a 3-day treatment with 4 days without medication, over a maximum period of 3 months.

APLONAL 5mg

The maximum potency, non-steroid anti-inflammatory, analgesic and antipyretic agent for ambulatory treatment.



FOR USE IN



DISTRIBUTION COUNTRIES



A non-steroid anti-inflammatory, analgesic and antipyretic agent for oral administration in dogs, equivalent to morphine in potency.

FORMULA

Each tablet contains: flunixin base (as meglumine) 5 mg
and excipients, q.s.p. 300 mg.

INDICATIONS

Pain and inflammation relief in dogs, irrespective of its origin,
especially musculo-skeletal.

ADMINISTRATION AND DOSAGE

1 tablet per 5 Kg weight every 24 hours, preferably with food, during a maximum of 3 consecutive days.
In chronic cases alternate a 3-day treatment with 4 days without medication, over a maximum period of 3 months.

ANESTHESICS

MYORELAXANTS

WWW.KONIGLAB.COM

KENSOL

Myorelaxant and anesthetic sedative.



FOR USE IN



DISTRIBUTION COUNTRIES



Injectable sedative, myorelaxant, analgesic and anesthetic agent for small and large animals.

FORMULA

Each ml contains: xylazine hydrochloride 20 mg;
excipients and vehicle q.s.

INDICATIONS

Sedative, myorelaxant, analgesic and anesthetic for use alone
or in association with preanesthetics or anesthetics
for contention maneuvers or major or minor surgery
in all domestic animals.

ADMINISTRATION AND DOSAGE

■ **Cattle:** 0.25-0.5 ml/100 kg, via IM or IV, for minor surgery, sedation or transportation purposes. Onset of action at 30 seconds for a duration of, approximately, 2 hours. Major surgery: 1 ml/100 kg, via IM or IV, with a 2 to 4 hour duration of effect. Kensol (0.5 ml/100 kg) can also be combined with Vetanarcol (10 ml/100 kg).

■ **Horses:** 3-10 ml/100 kg for sedation and light anesthesia, via SC, IM or IV. Duration of effect: between 15 and 30 minutes. For major surgery: combine 4 ml/100 kg Kensol (IM or IV) with an injection of 4-8 ml/100 kg Vetanarcol one hour later.

VETARNARCOL

Safe anesthetic. Cardiovascular stimulant. No effects on respiratory system.



FOR USE IN



DISTRIBUTION COUNTRIES



Dissociation acting general anesthetic for domestic and wild animals. Ideal for emergencies and traumatisms.

FORMULA

Each ml contains: ketamine (as hydrochloride) 50 mg, excipients and vehicle q.s.

INDICATIONS

Vetarnarcol is an anesthetic indicated for premedication, induction or maintenance use, alone or associated with major and minor tranquilizers, sedatives, narcotics, analgesics and other anesthetics for all kinds of domestic and wild animals.

ADMINISTRATION AND DOSAGE

■ **IV recommended guidance doses are:** horses: 3 to 8 mg/kg; cattle: 5 mg/kg; sheep and goats: 3 mg/kg; pigs: 5 mg/kg. ■ **IM recommended guidance doses are:** horses: 15 mg/kg; cattle: 15 mg/kg; sheep and goats: 10 to 20 mg/kg; pigs: 15 to 20 mg/kg.

It is recommended that Vetarnarcol be used associated with other drugs to obtain an anesthetic protocol adequate to each need, emphasizing its effects and minimizing its secondary adverse reactions. Vetarnarcol can be combined with xylazine (Kensol); diazepam, midazolam, acepromazine, barbiturics, propofol, narcotics, inhalatory agents and local anesthetics. The combination with other anesthetics and/or tranquilizers can reduce the required dose up to 50%, depending on the chosen protocol.

SMALL ANIMALS

ANTIBIOTICS

ANTI-INFLAMMATORY PRODUCTS

WWW.KONIGLAB.COM

BACTROPROFEN

Bacteriostatic and bactericide antibiotic that penetrates the infected core.



Oral administration antibiotic and steroid antiinflammatory agent for dogs and cats.

FORMULA

Each tablet contains: tetracycline hydrochloride 90 mg;
chloramphenicol 90 mg; base prednisolone 2 mg
and excipients q.s.p. 300 mg.

INDICATIONS

Treatment of general infectious processes with antibiotic,
antiinflammatory and analgesic action in dogs and cats.

ADMINISTRATION AND DOSAGE

1 tablet/10 kg weight every 12 hours. Duration of the treatment will depend on the evolution of the disease and is up to the criterion of the veterinarian. Recommended duration of the treatment is not less than 7 days.



FOR USE IN



DISTRIBUTION COUNTRIES



BTK 35

Cure and prevention of keratoconjunctivitis. Total acting bactericide, fungicide and viricide. Antiinflammatory.



Topical administration antibacterial and antiinflammatory agent for small and large animals.

FORMULA

Each 100 g of concentrate: argentic sulfadiazine 0,85 g%;
Oxytetracycline hydrochloride 5,00 g%; hydrocortisone acetate 0.29 g%; excipients q.s.f.

INDICATIONS

For the prevention and treatment of infectious keratoconjunctivitis in cattle, as well as for the prevention and treatment of bacterial inflammation and infection of traumatic and surgical wounds in horses.

ADMINISTRATION AND DOSAGE

Shake the container vigorously and spray locally over the affected area until a consistent and yellowish layer is formed. The container must be placed at some 20 to 30 cm distance from the area to be treated.

In cattle keratoconjunctivitis, apply once daily on both eyes during 5 days. In the case of wounds, clean thoroughly and apply twice a day.



FOR USE IN



DISTRIBUTION COUNTRIES



POLIDERM

Otic-dermic bactericide, fungicide and anti-inflammatory solution.



Antibiotic, anti-inflammatory and fungicidal solution for otic and dermatological use.

FORMULA

100 ml contain: thiabendazole 4 g; triamcinolone acetonide 0.1 g; neomicine base (as sulfate) 0.288 g and excipients q.s.

INDICATIONS

External otitis of bacterial and/or mycotic origin and localized dermatitis, with the same origin, in dogs and cats.

ADMINISTRATION AND DOSAGE

- **External otitis:** Clean the auditory canal carefully and instil 4/8 drops, twice daily.
- **Localized dermatitis:** apply on the affected part twice daily. In all cases, after the symptoms have totally disappeared, the treatment must be continued for 2/3 days more.



FOR USE IN



DISTRIBUTION COUNTRIES



ANTI-SEIZURE

WWW.KONIGLAB.COM

EPISOL

Sequential effect anticonvulsant and tranquilizer.



FOR USE IN



DISTRIBUTION COUNTRIES



Oral administration anticonvulsant and tranquilizer for dogs.

FORMULA

Each tablet contains: sodium diphenylhydantoinate 100 mg;
diazepam 5 mg and excipients q.s.p. 300 mg.

INDICATIONS

Control of central origin convulsions or epileptiform seizures.
Neuroregulator tranquilizer in cases of hyperexcitability,
aggressiveness and hypersexuality in dogs.

ADMINISTRATION AND DOSAGE

- **As anticonvulsant:** 1 tablet per 10 kg weight divided in two daily doses, during the first 20 days, then adjusting the supportive dose to the patient's clinical response.
- **As a tranquilizer:** 1/2 to 1 tablet per 10 kg weight divided in two daily doses.

CONTRACEPTIVES

WWW.KONIGLAB.COM

SINGESTAR COMPRIMIDOS

Minimum dose control of oestrus and pseudopregnancy. Feline dermatology.



FOR USE IN



DISTRIBUTION COUNTRIES



Contraceptive progestogen and for dermatological use in dogs and cats.

FORMULA

Each tablet contains: megestrol acetate 20 mg,
excipients q.s.p. 280 mg.

INDICATIONS

■ **For female dogs and cats:** prevention or annulment
of oestrus, treatment of estrogen dependent tumors. (*)

(*) **For dogs:** treatment of hypersexuality; perianal gland adenomas and prostate hypertrophy. **For cats:** treatment of hypersexuality; miliary dermatitis and eosinophilic granuloma.

ADMINISTRATION AND DOSAGE

■ **Bitches: For prevention or annulment of oestrus:** one tablet per 10 kg weight per day during 8 days (normal regime) or one tablet per 10 kg weight per day during 4 days and 1/4 tablet per 10 kg weight per day during 16 days (prolonged regime). **To delay or postpone oestrus:** 1/4 tablet per 10 kg weight per day during 30 to 40 days (normal regime) or 1/4 tablet per 10 kg weight per day during 30 to 40 days, then continuing with 1/4 tablet per 10 kg weight, twice a week, during a period not longer than 4 months (prolonged period). In the case of estrogen dependent tumors, one tablet per 10 kg weight per day during 5 days, then continuing with 1/2 tablet per 10 kg weight per day during 10 days or more. ■ **Female cats: For prevention or annulment of oestrus:** 1/2 tablet per day during 7 days, then continuing with 1/4 tablet, once or twice a week. **To delay or postpone oestrus:** 1/4 tablet during a period not longer than 2 months or 1/4 tablet once a week during a period not longer than 18 months. ■ **Dogs:** for hypersexuality: 1 or 2 tablets per 10 kg weight during 14 days (initial dose). In some cases, a maintenance dose of 1/2 tablet per 10 kg weight will be administered once or twice a week. **In perianal gland adenomas:** one tablet per 10 kg weight per day during 20 days or more. In prostate hypertrophy: One tablet per 10 kg weight per day, then continuing with 1/2 tablet per 10 kg weight twice a week. ■ **Cats: For hypersexuality:** 1/2 tablet per day during 7 days, continuing with 1/4 tablet per day during 14 days. A maintenance dose of 1/4 tablet may be indicated once or twice a week. **Miliary dermatitis:** 1/2 tablet per day during 7 days, continuing with 1/4 tablet twice a week until total cure is achieved. Eosinophilic granuloma: 1/2 tablet per day on alternate days up to initiation of regression, continuing with 1/2 tablet once a week until total regression.

SINGESTAR HI-JECT

Sterile, prefilled syringe. Atraumatic.



FOR USE IN



DISTRIBUTION COUNTRIES



Contraceptive progestogen in injectable suspension for female dogs and cats.

FORMULA

Each ml contains: medroxyprogesterone acetate 50 mg,
excipients and vehicle q.s.

INDICATIONS

Prevention or annulment of oestrus in bitches and female cats.

ADMINISTRATION AND DOSAGE

Apply 1 ml total per SC, only in the anestrus phase. To achieve an effective control in she-dogs, the treatment must be effected 20 to 30 days before the expected oestrus. In these animals, the treatment must be repeated every 4 to 6 months, according to the annual oestrus frequency.

■ **In female cats:** Administration must be carried out 3 to 6 weeks after the oestrus or a week after weaning of the kittens, both instances corresponding to the anestrus phase. It should be borne in mind that in a totally domestic environment, female cats may not present signs of the anestrus period. In these cases, the use of Singestar tablets is indicated.

SINGESTAR MP

Maximum effect contraceptive at a minimum dose.



FOR USE IN



DISTRIBUTION COUNTRIES



Contraceptive progestogen in injectable suspension for female dogs and cats.

FORMULA

Each ml contains: medroxyprogesterone acetate 50 mg,
excipients and vehicle q.s.

INDICATIONS

Prevention or annulment of oestrus in bitches and female cats.

ADMINISTRATION AND DOSAGE

Apply 1 ml total per SC, only in the anestrus phase. To achieve an effective control in bitches, the treatment must be effected 20 to 30 days before the expected oestrus. In these animals, the treatment must be repeated every 4 to 6 months, according to the annual oestrus frequency of oestrus.

■ **In female cats:** Administration must be carried out 3 to 6 weeks after the oestrus or a week after weaning of the kittens, both instances corresponding to the anestrus phase. It should be borne in mind that in a totally domestic environment, female cats may not present signs of the anestrus period. In these cases, the use of Singestar tablets is indicated.

ANTIDIARRHEIC PRODUCTS

WWW.KONIGLAB.COM

DIAFIN 2

Treatment of functional diarrhea in one single dose.



Injectable intestinal motility modulator antidiarrheal for large animals.

FORMULA

Each ml contains benzetimide hydrochloride 0.165 mg;
excipients and vehicle q.s.

INDICATIONS

Prevention and treatment of diarrheas in cattle originated
by nervous (penning, transfer, herding, transportation)
or functional (changes in feeding, grazing, weaning) conditions.
Symptomatic treatment of bacterial or viral complicated
diarrheas in cattle, sheep, horses and pigs.



FOR USE IN



DISTRIBUTION COUNTRIES



DIAFIN N

A single application stops diarrhea. Eliminates pathogen enterobacteria.



Injectable intestinal motility modulator antibiotic and antidiarrheal for small animals.

FORMULA

Each ml contains benzetimide hydrochloride 0.165 mg;
enrofloxacin 25 mg; excipients and vehicle q.s.

INDICATIONS

Treatment of complicated bacterial or viral diarrheas
in dogs and cats.

ADMINISTRATION AND DOSAGE

Apply 1 ml per each 10 kg weight, via SC or IM to both species.

In most cases, a single application is sufficient to cure diarrhea. Treatment can be repeated at 24 or 48 hours.



FOR USE IN



DISTRIBUTION COUNTRIES



EXTERNAL

ANTIPARASITICS

WWW.KONIGLAB.COM

DOMINAL COLLAR

External Antiparasitic Collar



FOR USE IN



DISTRIBUTION COUNTRIES



External antiparasitic for dogs and cats. The active ingredients are incorporated to the plastic matrix of the collar, as micro-particles that migrate to the surface continuously maintaining its efficacy in a sustained manner over a period of 3 months.

FORMULA

Contents per each 100 grams: Propoxur 12,00 grams; Imidacloprid 5,00 grams; Fenoxicarb, 1,00 grams, excipient q.s.

INDICATIONS

External antiparasitic destined to the prevention and treatment of the infestations by fleas tick and/or lice. The necessary dose is administered according to the weight of the collar and the size of the animal. It is recommended that the collar be changed every

ADMINISTRATION AND DOSAGE

Place the collar around the animal's neck and secure it after adjusting the size. Leave a space that allows the insertion of fingers, so as to avoid excessive tightness or too loose. Cut and discard the surplus piece. In the case of animals that are still growing, consider this situation, and allow a surplus to adjust the collar. Do not apply on puppies or kittens under two months old. Do not use on pregnant females. Remove the collar to bathe the animal and put it back on again when the animal is dry.

DOMINAL SOAP

Antiparasitic soap for dogs and cats.



Antiparasitic soap for dogs and cats with cosmetic action; indicated for the weekly bath.

FORMULA

Contents per each 100 grams: Imidacloprid 0,30 grams;
excipient q.s.

INDICATIONS

Treatment and prevention of the infestations by fleas and lice
in dogs and cats.

ADMINISTRATION AND DOSAGE

Wet the animal with warm water. Rub on to the body in the direction of hair growth and contrary to hair growth, until obtaining abundant foam so as to facilitate product contact and penetration. Allow to act for 3 minutes. Rinse completely and repeat the operation. Allow to dry naturally.



FOR USE IN



DISTRIBUTION COUNTRIES



DOMINAL MAX CATS

Antiparasitic with topical action and environmental effect for cats.



FOR USE IN



DISTRIBUTION COUNTRIES



Dominal MAX Cats is a topical application pulicicide solution for cats that incorporates the environmental effect. Dominal MAX Cats contains Imidacloprid and Fenoxicarb. Imidacloprid is characterized by its potent dumping power, responsible for the elimination of 100% of the fleas within 24 hours post-application. The Fenoxicarb provides residual power, due to its analogous action to that of the juvenile hormones of the flea, preventing the **maturing of eggs and larvae, eliminating same from the quarters without resorting to other additional products.**

FORMULA

Contents per each 100 mL: Imidacloprid 8,0 grams; Fenoxicarb 0,5 grams; excipient q.s.

INDICATIONS

Dominal MAX Cats is destined to the prevention and treatment of the infestations by fleas and lice.

ADMINISTRATION AND DOSAGE

Remove the spot-on from its pack. Maintain in a vertical position with the tip upwards and proceed to twist the point with your fingers and break it. Separate the animal's hair at the height of the base of the neck and apply the total contents of the spot-on directly upon

DOMINAL MAX DOGS

Antiparasitic with topical action and environmental effect for dogs.



FOR USE IN



DISTRIBUTION COUNTRIES



Dominal MAX Dogs is a topical application, pulicicide solution for dogs that incorporates environmental effect. **Dominal MAX Dogs** contains Imidacloprid, Permethrin and Fenoxycarb. Imidacloprid is characterized by its potent dumping power, responsible for the elimination of 100% of the fleas within the 24 hours post-application.. Permethrin provides the tickicide power. The Fenoxycarb furnishes residual power, due to its analogous action to that of the juvenile, hormones of the flea, preventing the eggs and larvae from maturing, eliminating same from the quarters without resorting to additional products.

FORMULA

Contents per each 100 mL: Permethrin 45 grams; Imidacloprid 10,0 grams; Fenoxycarb 0,5 grams; excipient q.s.

INDICATIONS

Prevention and treatment of infestations by fleas, ticks and lice. Repels flies and mosquitoes that transmit dirofilariasis (heart worm)

ADMINISTRATION AND DOSAGE

Remove the spot-on from its pack. Maintain in a vertical position with the tip upwards and proceed to twist the point with your fingers and break it. Separate the animal's hair at the height of the base of the neck and apply the total contents of the spot-on directly upon the skin. For dogs over 25 Kg. It is recommended that the application be made on two or three spots, separating the hair along the spine. Do not apply on cats.

DOMINAL POWDER

External antiparasitic powder. Effective against fleas, tick and lice.



External Antiparasitic for dogs and cats, that kills upon contact. It interrupts the cycle, preventing new re-infestations.

FORMULA

Contents for each 100 gr. : Propoxur 1,00 gram;
Imidacloprid 0,15 grams; excipient q.s.

INDICATIONS

It is indicated for the prevention and treatment of infestations by fleas, ticks and lice. Specially indicated for puppies and kittens owing to its low toxicity.

ADMINISTRATION AND DOSAGE

Sprinkle softly along the spine, groin and axilla. Spread with a massage on the hair coating, in a direction contrary to the hair growth. It is recommended that the procedure be repeated once



FOR USE IN



DISTRIBUTION COUNTRIES



DOMINAL SHAMPOO

Antiparasitic Shampoo. Hygiene and Cleanliness of the skin in dogs and cats.



External Antiparasitic, pulicide, tickicide and pediculicides for dogs and cats, that kills upon contact.

FORMULA

Contents for each 100 mL: Propoxur, 0,165 gr; Imidacloprid, 0,150 gr; excipient q.s.

INDICATIONS

Indicated for prevention and treatment of infestations by fleas, tick and lice. It possesses an excellent cleansing power, leaving the skin and hair hygienic.

ADMINISTRATION AND DOSAGE

Wet the animal's hair with warm water and apply the shampoo; create abundant foam and rinse, Repeat the procedure and allow the foam to act for a few minutes before the last rinse. It is recommendable that the baths be repeated every 7 days but this frequency may vary according the veterinarian's suggestion. Do not apply on kittens or puppies under two months old.



FOR USE IN



DISTRIBUTION COUNTRIES



DOMINAL CATS

Lethal Impact against fleas.



FOR USE IN



DISTRIBUTION COUNTRIES



Dominal Cats is a pulicicide solution for cats.

Dominal Cats contains Imidacloprid. Imidacloprid is characterized by its potent dumping power, responsible for the elimination of 100% of the fleas within 24 hours as from post-application.

Its rapid and effective adulticide and larvicide action leads to the flea's cycle being cut since they are eliminated before they begin to lay eggs. Dominal Cats protects against reinfestations by fleas.

FORMULA

Contents per each 100 mL: Imidacloprid 8 grams;
excipient q.s.

INDICATIONS

Dominal Cats is destined to the prevention and treatment of infestations by fleas.

ADMINISTRATION AND DOSAGE

Remove the spot-on from its pack. Maintain in a vertical position with the tip upwards and proceed to twist the point with your fingers and break it. Separate the animal's hair at the height of the base of the neck and apply the total contents of the spot-on directly upon

DOMINAL DOGS

Lethal Impact against fleas and ticks.



+

FOR USE IN



+

DISTRIBUTION COUNTRIES



Dominal Dogs is a topical pulicide and tickicide solution for dogs.

Dominal Dogs contains Imidacloprid and Permethrin. Imidacloprid is characterized by its potent killing power responsible for the elimination of 100% of the fleas within 24 hours post-application.

Its rapid and effective adulticide and larvicide action leads to the life cycle of fleas being cut since they are eliminated before they start to lay eggs. Permethrin furnishes the tickicide power. Dominal Dogs prevents reinfestations by fleas and ticks.

FORMULA

Contents per each 100 mL: Permethrin 45 grams; Imidacloprid 10 grams; excipient q.s.

INDICATIONS

Treatment and prevention of infestations by fleas and ticks in dogs. Adulticide and larvicide.

ADMINISTRATION AND DOSAGE

Remove the spot-on from its pack. Maintain in a vertical position with the tip upwards and proceed to twist the point with your fingers and break it. Separate the animal's hair at the height of the base of the neck and apply the total contents of the spot-on directly upon the skin. For dogs over 25 Kg. It is recommended that the application be made on two or three spots, separating the hair along the spine. Do not apply on cats.

DOMINAL SPRAY

Kills fleas, ticks and lice on dogs. The only product with genuine four-way effect against adult fleas, eggs, larvae and pupae.



The state-of-the-art in aerosol spray. Kills fleas, ticks and lice on dogs.

FORMULA

100 ml contains: imidacloprid 0.15 g; permethrin 1 g;
fenoxycarb 0.06 g; piperonyl butoxide 3 g and excipients q.s.

INDICATIONS

Treatment and prevention of infestation with fleas,
ticks and lice, in dogs.

ADMINISTRATION AND DOSAGE

Spray on animal's body surface. Make sure that the liquid comes into contact with skin.
Repeat the application every 30 days.



FOR USE IN



DISTRIBUTION COUNTRIES



FAUNA SOAP

Eliminates fleas, ticks and mange mites.



FOR USE IN



DISTRIBUTION COUNTRIES



Ectoparasiticide and hygiene soap for dogs and cats.

FORMULA

100 g contains: amitraz 0.125g; sulfur 2 g and excipients q.s.

INDICATIONS

Weekly bath in dogs and cats, specially in case of infestation with fleas, ticks, lice, and sarcoptic, notoedric and demodectic mange.

ADMINISTRATION AND DOSAGE

Wet hair with abundant lukewarm water and rub vigorously allowing the lather to act for 3/5 minutes. Repeat the operation. In case of infestation with fleas or ticks, repeat weekly. For treatment of demodectic, bathe once a week during a minimum period of two months. In sarcoptic and notoedric mange, an interval of 4 days between baths is recommended and the duration of the treatment depends on the evolution of the condition.

TEA 327 AEROSOL

Kills fleas, ticks and lice. Deodorant.



Aerosol Spray for fleas, ticks and lice control, with deodorant action, for small animals.

FORMULA

100 g of concentrate contains: propoxur 2.26 g and excipients q.s. Propellant: butane-propane.

INDICATIONS

Infestation with fleas, ticks and lice, in dogs and cats.
Treatment of seborreic dermatitis

ADMINISTRATION AND DOSAGE

Spray against the natural lay of the hair, on all the animal's body, keeping the container at 10 to 15 cm distance.
Use also recommended in kennels and sleeping quarters.



FOR USE IN



DISTRIBUTION COUNTRIES



TEA 327 CAT COLLAR

Fleas and lice control collars. Effective for 5 months.



Fleas, and lice control collars for cats.

FORMULA

100 g contains: propoxur 10 g and excipients q.s.

INDICATIONS

Infestation with fleas and lice, in cats.

ADMINISTRATION AND DOSAGE

Place the collar on the animal's neck, being careful that it is neither too tight nor loose, allowing for two fingers to pass under it. Cut off excess material. Change the collar every 5 months.



FOR USE IN



DISTRIBUTION COUNTRIES



TEA 327 DOG COLLAR

Fleas, ticks and lice control collars. Effective for 5 months.



Fleas, ticks and lice control collars for dogs.

FORMULA

100 g contains: propoxur 10 g and excipients q.s.

INDICATIONS

Infestation with fleas, ticks and lice, in dogs.

ADMINISTRATION AND DOSAGE

Place the collar on the animal's neck, being careful that it is neither too tight nor loose, allowing for two fingers to pass under it. Cut off excess material. Change the collar every 5 months.



FOR USE IN



DISTRIBUTION COUNTRIES



TEA 327 POWDER

Dry Control for fleas and ticks. Ideal for in-between baths.



Powder for fleas, ticks and lice control in small animals.

FORMULA

100 g contains: propoxur 1 g and excipients q.s.

INDICATIONS

Infestation with fleas, ticks and lice, in dogs and cats, especially puppies.

ADMINISTRATION AND DOSAGE

Apply on the animal's body every 7 days. Use recommended in kennels and sleeping quarters.



FOR USE IN



DISTRIBUTION COUNTRIES



TEA DERMASOFT

Dermocompatible rinse for fleas and ticks control. Non allergenic.



Rinse for fleas, ticks and lice control with cosmetic properties, for dogs.

FORMULA

100 ml contains: DDVP 2.8 g, cypermethrin 0.8 g
and excipients q.s.

INDICATIONS

Ectoparasitic infestation in dogs.

ADMINISTRATION AND DOSAGE

Dilute 15 ml in one litre of water. After bathing apply the resulting emulsion over all the animal's body using a sponge, working against the natural lay of the hair. Do not rinse and allow to dry naturally. Repeat every 7 days.



FOR USE IN



DISTRIBUTION COUNTRIES



TEA LARVOX

Environmental Aerosol Spray which totally cuts the flea cycle. The only one which acts against resistant strains.



Environmental aerosol spray control for fleas and ticks with action against eggs, larvae and adults.

FORMULA

100 g of aerosol spray contains natural pyrethrins 0.1 g,
methoprene 0.15 g, propoxur 1 g, DDVP 0.47 g,
piperonyl butoxide 0.4 g excipients and propellant q.s.

INDICATIONS

Eliminates fleas and ticks from the environment.

ADMINISTRATION AND DOSAGE

Apply in kennels, carpets, cushions, floors, corners and skirting boards in places where dogs and cats usually go.
One single fumigation is effective for 120 days.



FOR USE IN



DISTRIBUTION COUNTRIES



TEA SUPER SHAMPOO

Hair beauty and sheen. Kills fleas and ticks.



Ectoparasiticide and grooming shampoo for dogs.

FORMULA

100 ml contains: natural pyrethrins 0.1 g;
piperonyl butoxide 1 g, N-octyl bicycloheptene
dicarboximide 0.5 g and excipients q.s.

INDICATIONS

Kills fleas and ticks and hair care in dogs.

ADMINISTRATION AND DOSAGE

Wet hair with lukewarm water. Apply shampoo and massage until you create an abundant lather, rinse and repeat the operation. In the second application allow to act a few minutes before rinsing. Repeat bath every 7 days.



FOR USE IN



DISTRIBUTION COUNTRIES



INTERNAL

ANTIPARASITICS

WWW.KONIGLAB.COM

BASKEN DOBLE 20

Total spectrum, single dose antiparasitic agent, with no side effects.



FOR USE IN



DISTRIBUTION COUNTRIES



Broad spectrum anthelmintic for oral administration to dogs.

FORMULA

Each tablet contains: pyrantel pamoate 72.08 mg;
oxantel pamoate 69.90 mg; praziquantel 25 mg
and excipients, q.s.p. 550 mg.

INDICATIONS

Treatment of parasitosis caused by nematodes (*Toxocara canis*; *Ancylostoma caninum*; *Uncinaria stenocephala*; *Trichuris vulpis*) and cestodes (*Echinococcus granulosus*, *Dipylidium caninum* and taenias in general) in dogs, as from 30 days old. (*)

(*) It possesses larvae and adult elimination activity. It may be administered during pregnancy without any risk and has no effect on fertility in either males or females.

ADMINISTRATION AND DOSAGE

1 tablet, in one single dose, for every 5 Kg weight. Repeat after 15 days. In case of severe trichuriasis, a treatment of 3 consecutive days is recommended. There is no need for previous or subsequent fasting. It may be administered directly or mixed with food.

BASKEN DOBLE 40

Total spectrum, single dose antiparasitic agent, with no side effects.



FOR USE IN



DISTRIBUTION COUNTRIES



Broad spectrum anthelmintic for oral administration to dogs.

FORMULA

Each tablet contains: pyrantel pamoate 144.12 mg;
oxantel pamoate 139.80 mg; praziquantel 50 mg
and excipients, q.s.p. 1.100 mg.

INDICATIONS

Treatment of parasitosis by nematodes (*Toxocara canis*;
Ancylostoma caninum; *Uncinaria stenocephala*; *Trichuris vulpis*)
and cestodes (*Echinococcus granulosus*, *Dipylidium caninum*
and taenias in general) in dogs, as from 30 days old. (*)

(*) It possesses larvae and adult elimination activity. It may be administered during pregnancy without any risk and has no effect on fertility in either males or females.

ADMINISTRATION AND DOSAGE

1 tablet, in one single dose, for every 10 Kg weight. Repeat after 15 days. In case of severe trichuriasis, a treatment of 3 consecutive days is recommended. There is no need for previous or subsequent fasting. It may be administered directly or mixed with food.

BASKEN DOBLE 60

Total spectrum, single dose antiparasitic agent, with no side effects.



FOR USE IN



DISTRIBUTION COUNTRIES



Broad spectrum anthelmintic for oral administration to dogs.

FORMULA

Each tablet contains: pyrantel pamoate 288.40 mg;
oxantel pamoate 279.60 mg; praziquantel 100 mg
and excipients, q.s.p. 1.300 mg.

INDICATIONS

Treatment of parasitosis by nematodes (*Toxocara canis*;
Ancylostoma caninum; *Uncinaria stenocephala*; *Trichuris vulpis*)
and cestodes (*Echinococcus granulosus*, *Dipylidium caninum*
and taenias in general) in dogs, as from 30 days old. (*)

(*) It possesses larvae and adult elimination activity. It may be administered during pregnancy without any risk and has no effect on fertility in either males or females.

ADMINISTRATION AND DOSAGE

1 tablet, in one single dose, for every 20 Kg weight. Repeat after 15 days. In case of severe trichuriasis, a treatment of 3 consecutive days is recommended. There is no need for previous or subsequent fasting. It may be administered directly or mixed with food.

BASKEN PLUS DOGS

Total spectrum, single dose antiparasitic agent, with no side effects.



FOR USE IN



DISTRIBUTION COUNTRIES



Broad spectrum anthelmintic for oral administration to small dogs.

FORMULA

Each tablet contains: pyrantel pamoate 36.03 mg;
oxantel pamoate 34.95 mg; praziquantel 6.24mg
and excipients, q.s.p. 310 mg.

INDICATIONS

Treatment of parasitosis caused by nematodes (*Toxocara canis*;
Ancylostoma caninum; *Uncinaria stenocephala* and *Trichuris*
vulpis) and cestodes (*Dipylidium caninum* and taenias
in general) in dogs, as from 30 days old. (*)

(*) It possesses larvae and adult elimination activity. It may be administered to females at any stage during pregnancy without any risk. It has no effect on fertility in either males or females.

ADMINISTRATION AND DOSAGE

1 tablet, in one single dose, for every 2,5 Kg weight. Repeat after 15 days. In case of severe trichuriasis, a treatment of 3 consecutive days is recommended. There is no need for previous or subsequent fasting. It may be administered directly or mixed with food.

BASKEN PLUS CATS

Total spectrum, single dose antiparasitic agent, with no side effects.



FOR USE IN



DISTRIBUTION COUNTRIES



Broad spectrum anthelmintic for oral administration to cats.

FORMULA

Each tablet contains: pyrantel pamoate 36.03 mg;
oxantel pamoate 34.95 mg; praziquantel 6.24mg
and excipients, q.s.p. 310 mg.

INDICATIONS

Treatment of parasitosis caused by nematodes (*Toxocara cati*; *Toxascaris leonina*; *Ancylostoma tubaeforme*; *Uncinaria stenocephala* and *Trichuris vulpis*) and cestodes (*Dipylidium caninum* and taenias in general) in cats, as from 30 days old. (*)

(*) It possesses larvae and adult elimination activity. It may be administered to females at any stage during pregnancy without any risk. It has no effect on fertility in either males or females.

ADMINISTRATION AND DOSAGE

1 tablet, in one single dose, for every 2,5 Kg weight. Repeat after 15 days. In case of severe trichuriasis, a treatment of 3 consecutive days is recommended. There is no need for previous or subsequent fasting. It may be administered directly or mixed with food.

BASKEN SUSPENSION

Total spectrum, single dose antiparasitic agent, with no side effects.



+

FOR USE IN



+

DISTRIBUTION COUNTRIES



Nematocide anthelmintic for pups and kittens.

FORMULA

Each ml contains: pyrantel pamoate 14.5 mg;
oxantel pamoate 14 mg and excipients, q.s.

INDICATIONS

Treatment of parasitosis caused by *Toxocara canis*, *Toxocara cati*,
Toxascaris leonina; *Ancylostoma* spp; *Uncinaria stenocephala*
and *Trichuris vulpis* in pups and kittens. It is also effective
in killing larvae and adults.

ADMINISTRATION AND DOSAGE

Administer 1 ml/kg weight, starting when the animal is 15 days old. Repeat every fortnight until the vaccination program is complete or repeat after 15 days and continue with the Basken tablets, adhering to the same scheme. In case of severe trichuriasis, a treatment of 3 consecutive days is recommended. Basken may be administered in females during any stage of pregnancy and during lactation, with offspring of more than 15 days. Administer directly using the attached dosing syringe, or mixed with a small portion of food. There is no need for previous or subsequent fasting. Basken does not affect fertility in either males or females.

CESTODAN 50

Total taenicide, with broad spectrum, for dogs and cats.



FOR USE IN



DISTRIBUTION COUNTRIES



Broad spectrum cestodicide for oral administration, in dogs and cats.

FORMULA

Each tablet contains: praziquantel 50 mg and excipients,
q.s.p. 525 mg.

INDICATIONS

Parasitosis caused by *Echinococcus granulosus*, *Dipylidium*
caninum and taenias in general in dogs and cats.

ADMINISTRATION AND DOSAGE

Administer 1 tablet per 10 Kg of weight in one single dose, directly in the mouth or mixed with food.
There is no need for previous fasting. It may be administered to pregnant dogs and cats. In areas of endemic
hydatidosis repeat treatment every 45 days.

CESTODAN INYECTABLE

Exclusive total taenicide for parenteral use.



Injectable, broad spectrum, painless cestodicide for dogs and cats.

FORMULA

Each ml contains: praziquantel 55 mg; excipients and vehicle q.s.

INDICATIONS

Parasitosis caused by *Echinococcus granulosus*, *Dipylidium caninum* and taenias in general in dogs and cats.

ADMINISTRATION AND DOSAGE

Apply 1 ml per every 10 Kg weight, exclusively by subcutaneous injection. In animals over 30 Kg weight, apply a maximum dose of 3 ml. It may be administered to pregnant dogs and cats. In areas of endemic hydatidosis repeat the treatment every 45 days.



FOR USE IN



DISTRIBUTION COUNTRIES



CESTODAN ORAL

Total taenicide, with broad spectrum, for dogs and cats.



FOR USE IN



DISTRIBUTION COUNTRIES



Broad spectrum, cestodicide for oral administration in dogs and cats.

FORMULA

One tablet contains: praziquantel 25 mg
and excipients q.s.p.

INDICATIONS

Parasitosis caused by *Echinococcus granulosus*, *Dipylidium caninum* and taenias in general, in dogs and cats.

ADMINISTRATION AND DOSAGE

One tablet every 5 Kg weight, as a single dose, directly in the mouth or mixed with food. No prior fasting required. It may be administered to pregnant dogs and cats. In areas of endemic hydatidosis repeat the treatment every 45 days.

VERMIPLEX

Internal Antiparasitic. Immunomodulator.



FOR USE IN



DISTRIBUTION COUNTRIES



Broad spectrum nematocide anthelmintic for dogs and cats.

FORMULA

Each tablet contains: thiabendazole 200 mg,
levamisole hydrochloride 6 mg, excipients q.s.p. 525 mg.

INDICATIONS

Treatment of parasitosis caused by nematodes (*Toxocara canis*,
Toxocara cati; *Toxascaris leonina*; *Ancylostoma* spp; *Uncinaria*
stenocephala ; *Trichuris vulpis*) in dogs and cats.

ADMINISTRATION AND DOSAGE

One tablet every 4 Kg of weight, fasting. Maximum dose: 6 tablets. Repeat after 15/20 days. May be administered as from 3 weeks old.

BASKEN 4 APR

Internal Antiparasitic for Dogs



FOR USE IN



DISTRIBUTION COUNTRIES



Wide spectrum Internal antiparasitic possessing action against resistant parasites.

FORMULA

Contents per each 100 grams: Pyrntel Pamoate 24,17 grams;
Oxantel Pamoate 23,33 grams;
Oxfendazol 18,83 grams; Praziquantel 8,33 grams; excipients
c.s.

INDICATIONS

Treatment of parasitoses that affect dogs, caused by nematodes (Toxocara canis, Ancylostoma caninum, Uncinaria stenocephala, Trichuris vulpiscanis); cestodes (Echinococcus granulosus, Dipilydium caninum and tapeworms in general) and Giardias spp.

ADMINISTRATION AND DOSAGE

- Administer 1- 150 mg tablet for each 2,5 Kg. of live weight.
 - Administer 1-300 mg tablet for each 5 Kg. of live weight.
 - Administer 1- 600 mg tablet for each 10 Kg. of live weight.
 - Administer 1-1200 mg tablet for each 20 Kg. of live weight.
 - Dose of actives per Kg. of weight: Pyrntel Pamoate 14,5 mg/Kg. (equivalent to 5 mg/Kg. de Pyrntel); Oxantel Pamoate 14,0 mg/Kg. (equivalent a 5 mg/Kg. of Oxantel); Praziquantel 5 mg/Kg.; Oxfendazol 11,3 mg/kg.
- Do not administer to Cats, puppies under 3 weeks old, females in the first third of their pregnancy term or breast-feeding with puppies under 21 days old.

CALCIFYING PRODUCTS

WWW.KONIGLAB.COM

EUTAZUM

High absorption calcium.



Mineral supplement, in oral suspension, for small animals.

FORMULA

100 ml contains: Calcium carbonate 10 g; flavour
and excipients q.s.

INDICATIONS

Prevention of disorders associated with the calcium demand
increase during nursing and growth stages.
Treatment of generalised fibrous osteodystrophy, nutritional
secondary hyperparathyroidism and hypocalcemia.

ADMINISTRATION AND DOSAGE

As a prevention, administer one measure/5 kg weight/day. As a cure administer 2 measures/5 Kg weight/day.
It may be given directly in the mouth or mixed with food.



FOR USE IN



DISTRIBUTION COUNTRIES



SMALL ANIMALS

PHYSIOLOGICAL CICATRIZATION PRODUCTS

WWW.KONIGLAB.COM

KURADERM PLATA

Physiological Healing Agent. Keratoplastic. No formation of crust. Accidental or surgical wounds.



Antimicrobial; healing; astringent and high adherence hemostatic agent, for topical use in small and large animals.

FORMULA

100 g of concentrate contains: Argentic sulfadiazine 1.25 g;
neomycin base (as sulfate) 5.4 g; aluminum 8.33 g
and excipients q.s. Propellant: Butane-propane.

INDICATIONS

Prevention and treatment of infectious processes in wounds.
Hemostatic effect. Healing of wounds, eschars, burns and ulcers.

ADMINISTRATION AND DOSAGE

Shake before applying. After cleansing the area to be treated, spray a thin layer on same from a distance of approximately 10 cm ensuring penetration inside the wound. Repeat once a day until the wound heals.



FOR USE IN



DISTRIBUTION COUNTRIES



BACTROVET PLATA AM PET

High adherence Hemostatic, Healing and Antimyasic Agent.



Healing, antimyasic, antimicrobial and high adherence hemostatic agent for topical use in small animals.

FORMULA

100 g of concentrate contains: Argentic sulfadiazine 0.1 g;
aluminum 5 g; DDVP 1.6 g; cypermethrin 0.4 g
and excipients q.s.

INDICATIONS

Prevention of bleeding, infection and myiasis in traumatic
and surgical wounds. Healing treatment of ulcerous skin myiasis.
With larvicide and healing stimulant effect. Use in small animals.

ADMINISTRATION AND DOSAGE

Shake the container before applying. Spray the wound or myiasis abundantly, from a distance of 10 cm.
Apply once a day until cure.



FOR USE IN



DISTRIBUTION COUNTRIES



HORMONAL PRODUCTS

WWW.KONIGLAB.COM

ECP ESTRADIOL

Minimum dose and long acting estrogen.



FOR USE IN



DISTRIBUTION COUNTRIES



Injectable hormonal agent for small and large animals.

FORMULA

Each 100 ml contains: estradiol cypionate 0.2 g;
excipients and vehicle q.s.

INDICATIONS

Cows and heifers: Anestrus without follicular cyst; retention of the corpus luteum in milk cows; expulsion of placental residues or mummified foetus.

Mares, ewes and sows: Anestrus.

ADMINISTRATION AND DOSAGE (IM Injection only)

Cows: anestrus: 1.5 to 2.5 ml total dose; corpus luteum persistence: 2.0 ml total dose; amnion hydrops: 3.0 to 4.0 ml total dose; pyometra, placenta retention and expulsion of mummified foetus: 5.0 ml total dose.

Heifers: Anestrus: 1.5 ml total dose;

Mares: anestrus: 2.5 to 5.0 ml total dose;

Ewes and sows: Anestrus: 0.25 to 0.5 ml total dose.

iTHANK YOU!



CONSTRUCTING KNOWLEDGE
